

One-loop Energy Corrections in the Nonabelian Gauge Field Theory in 2+1 Dimensions

V. Ch. Zhukovsky, N. A. Peskov*
*Physical Faculty, Moscow State University,
119899, Moscow, Russia*

Abstract

Exact constant solutions of field equations in the classical nonabelian $SU(2)$ field theory with the Chern-Simons topological mass in $2 + 1$ -dimensional space-time have been obtained. One-loop contributions of boson and fermion fluctuations to the gauge field energy have been calculated.

arXiv:hep-th/9812221v2 19 Jan 2000

*E-mail addresses: th180@phys.msu.su and peskov_nick@mail.ru.

It is well known that a topological term, the so-called Chern-Simons term, which plays the role of the gauge field mass, can be introduced in odd-dimensional gauge theories [1, 2].

As an example of such theories, we consider the three-dimensional topologically massive $SU(2)$ -gluodynamics described by the Lagrangian [1]

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu} - \frac{\theta}{4}\varepsilon^{\mu\nu\alpha} \left(F_{\mu\nu}^a A_\alpha^a - \frac{g}{3}\varepsilon^{abc} A_\mu^a A_\nu^b A_\alpha^c \right). \quad (1)$$

Here $\mu, \nu = 0, 1, 2$; $a = 1, 2, 3$, potentials $A_\mu \equiv \tau^a A_\mu^a/2$, where τ^a are Pauli matrices in color space, the color field tensor is given by $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g\varepsilon^{abc} A_\mu^b A_\nu^c$. The coefficient θ in front of the last term in (1) (Chern-Simons term) is the Chern-Simons (CS) mass of the gauge field.

In this case the field equations take the form

$$D_\mu^{ac} F^{a\mu\beta} - \frac{\theta}{2}\varepsilon^{\beta\mu\nu} F_{\mu\nu}^c = 0, \quad (2)$$

where $D_\mu^{ac} = \delta^{ac}\partial_\mu - g\varepsilon^{abc}A_\mu^b$. It is easily verified that these equations are satisfied by the following constant nonabelian potentials

$$A^{a\mu} = \frac{\theta}{2g}\delta^{a\mu}\chi_{\lambda\omega}^{(a)}, \quad (3)$$

with normalized constant vector $\chi_{\lambda\omega}^{(a)} = (\lambda i, \lambda\omega i, \omega)$, where $\lambda = \pm 1$ and $\omega = \pm 1$. Note that λ and ω take its values independently. The Kronecker delta $\delta^{a\mu}$ in (3) implies that directions 1, 2, 3 in the color space correspond to directions 1, 2, 0 in the Minkowski 2 + 1 space-time, respectively. The following relations are easily verified:

$$F_{\mu\nu}^a = -\frac{\theta}{2}\varepsilon_{\mu\nu\alpha}A^{a\alpha}, \quad A^{a\mu}A_\mu^b = \frac{\theta^2}{4g^2}\delta^{ab}, \quad A^{a\mu}A^{a\nu} = \frac{\theta^2}{4g^2}g^{\mu\nu}.$$

The gauge field energy density E corresponds to the $\mathcal{T}^{00}$ component of the symmetric energy-momentum tensor $\mathcal{T}^{\mu\nu}$ obtained by the standard method [3]

$$\mathcal{T}^{\mu\nu} = T^{\mu\nu} + \partial_\alpha T^{\alpha\mu\nu} = {}^*F^{a\mu}{}^*F^{a\nu} - \frac{1}{2}g^{\mu\nu}{}^*F_\alpha^a{}^*F^{a\alpha}.$$

Here $T^{\mu\nu}$ is the canonical energy-momentum tensor, $T_{\alpha\mu\nu} = -A_\mu^a \left(F_{\alpha\nu}^a + \frac{1}{2}\theta\varepsilon_{\alpha\nu\beta}A^{a\beta} \right)$, and ${}^*F_\mu^a = \frac{1}{2}\varepsilon_{\mu\alpha\beta}F^{a\alpha\beta}$, the dual field tensor, is in fact a vector in 2 + 1-dimensional space-time. Taking the solution (3) into account we can now represent $\mathcal{T}^{\mu\nu}$ in the form

$$\mathcal{T}^{\mu\nu} = -\frac{\theta^4}{32g^2}g^{\mu\nu}.$$

The negative sign of the energy density $E = \mathcal{T}^{00}$ means that, for the given value of $\theta \neq 0$, the energy of this solution is lower than that of the trivial vacuum solution $A^\mu = 0$, obtained in the perturbation theory.

Next, we consider the one-loop effective potential

$$V_{\text{eff}} = E + \mathcal{E}_G^{(1)} + \mathcal{E}_Q^{(1)}$$

with consideration for both gluon $\mathcal{E}_G^{(1)}$ and quark $\mathcal{E}_Q^{(1)}$ loops. We start with the gluon contribution $\mathcal{E}_G^{(1)}$. The total gauge field $\mathcal{A}^{a\mu}$ is represented as the sum of the constant classical background field (3) and quantum fluctuations $a^{a\mu}$, i.e.

$$\mathcal{A}^{a\mu} = A^{a\mu} + a^{a\mu}.$$

Introducing the gauge-fixing term

$$\mathcal{L}_{\text{gf}} = -\frac{1}{2\xi} \left(\nabla_\mu^{ab} a^{b\mu} \right)^2,$$

where $\nabla_\mu^{ab} = \delta^{ab} \partial_\mu + g \varepsilon^{abc} A_\mu^c$ is the background covariant derivative, we obtain the fluctuation Lagrangian for the gauge $\xi = 1$ in the form

$$\mathcal{L}(a_\mu^b | A_\mu^b) = \mathcal{L}(a_\mu^b) - \frac{1}{2} (\partial_\mu a^{b\mu})^2 - \frac{\theta^2}{4} (a^{b\mu})^2 + g \varepsilon^{abc} A_\mu^a f^{b\mu\nu} a_\nu^c - g \varepsilon^{abc} A_\mu^c a^{b\mu} \partial_\nu a^{a\nu}.$$

Here, the first term in r.h.s. is the same Lagrangian as in (1), though defined in terms of the field fluctuations a_μ^b and the corresponding tensor $f_{\mu\nu}^b$.

The energy spectrum of gluon fluctuation modes ε_i , corresponding to *all* degrees of freedom (physical and unphysical) that remain after gauge fixing, is determined by the equation

$$\det \left[\delta^{ac} g^{\mu\nu} \left(p^2 + \frac{1}{2} \theta^2 \right) - i \theta \delta^{ac} \varepsilon^{\alpha\mu\nu} p_\alpha - 2i g \varepsilon^{abc} \left(A^{b\nu} p^\mu + A^{b\mu} p^\nu - g^{\mu\nu} A_\alpha^b p^\alpha \right) \right] = 0.$$

Nine solutions of the above equation are as follows:

$$\varepsilon_i^2 = \mathbf{p}^2 + \lambda_i \frac{\theta^2}{2},$$

where $\lambda_{1,2} = 3 \pm 2\sqrt{2}$, $\lambda_{3,4} = 5 \pm 2\sqrt{6}$, and the other five degenerate spectrum branches with $\lambda_{5,\dots,9} = -1$, corresponding to tachyonic modes, are unstable.

The one-loop energy correction due to gauge field fluctuations around the background is given by the expression

$$\mathcal{E}_G^{(1)} = \frac{1}{2} \sum_{\mathbf{p}, i} \varepsilon_i(\mathbf{p}) = \frac{1}{2} \sum_{\mathbf{p}, i} [\mathbf{p}^2 + \lambda_i \theta^2 / 2]^{1/2}. \quad (4)$$

Consideration for tachyonic modes contribution to expression (4) results in the appearance of the imaginary part of the energy correction $\mathcal{E}_G^{(1)}$, defined as follows (see, e.g., [3]):

$$\text{Im } \mathcal{E}_G^{(1)} = -\frac{1}{2} \sum_{\varepsilon_i^2 < 0} |\varepsilon_i^2(\mathbf{p})|^{1/2}.$$

The imaginary part is finite and its calculation results in the following expression (with due regard for degeneracy)

$$\text{Im } \mathcal{E}_G^{(1)} = -\frac{5}{2(2\pi)^2} \int_{|\mathbf{p}|=0}^{\theta/\sqrt{2}} \sqrt{\frac{\theta^2}{2} - \mathbf{p}^2} d\mathbf{p} = -\frac{5}{24\sqrt{2}\pi} |\theta|^3.$$

The real part of the energy correction corresponding to $\varepsilon_i^2 > 0$ contribution to (4) is divergent. The appropriate application of the dimensional regularization procedure yields:

$$\text{Re } \mathcal{E}_G^{(1)reg} = \frac{1}{2} \mu^{2\epsilon} \sum_{\varepsilon_i^2 > 0} [\mathbf{p}^2 + \lambda_i \theta^2 / 2]^{1/2-\epsilon}, \quad (5)$$

where μ is an arbitrary scale mass parameter. We can now introduce the proper time representation for this expression, and using the regulator $\epsilon > \frac{1}{2}$, we find

$$\text{Re } \mathcal{E}_G^{(1)reg} = \frac{\mu^{2\epsilon}}{2(2\pi)^2 \Gamma(\epsilon - 1/2)} \int_0^\infty \frac{ds}{s^{3/2-\epsilon}} \left\{ \int d\mathbf{p} \sum_{i=1}^4 e^{-s(\mathbf{p}^2 + \lambda_i \theta^2 / 2)} + 5 \int_{|\mathbf{p}|=\theta/\sqrt{2}}^\infty d\mathbf{p} e^{-s(\mathbf{p}^2 - \theta^2 / 2)} \right\},$$

and this, upon integration over the two-dimensional momentum, yields

$$\text{Re } \mathcal{E}_G^{(1)reg} = \frac{\mu^{2\epsilon}}{8\pi \Gamma(\epsilon - 1/2)} \int_0^\infty \frac{ds}{s^{5/2-\epsilon}} \left\{ 5 + \sum_{i=1}^4 e^{-s\lambda_i \theta^2 / 2} \right\}.$$

The (ultraviolet) divergent at $s \rightarrow 0$ term in $\text{Re } \mathcal{E}_G^{(1)reg}$ is determined by the expression

$$\text{Re } \mathcal{E}_G^{(1)div} = \frac{\mu^{2\epsilon}}{8\pi \Gamma(\epsilon - 1/2)} \int_0^\infty \frac{ds}{s^{5/2-\epsilon}} \left\{ 5 + \sum_{i=1}^4 \left(1 - s\lambda_i \frac{\theta^2}{2} \right) \right\}. \quad (6)$$

The difference $\text{Re } \mathcal{E}_G^{(1)ren} \equiv \text{Re } \mathcal{E}_G^{(1)reg} - \text{Re } \mathcal{E}_G^{(1)div}$ is finite in the limit $\epsilon \rightarrow 0$ and its calculation can be reduced to Γ -function integrals. However, the two results, for imaginary and real parts, still include contributions of the unphysical modes introduced by the gauge-fixing term. In order to get rid of these unphysical contributions, the ghost fields η^a and $\bar{\eta}^a$ with the Lagrangian

$$\mathcal{L}_{gh} = (\nabla_\mu^{ab} \bar{\eta}^b) (D^{ac\mu} \eta^c)$$

should be introduced. Here $D_\mu^{ab} = \delta^{ab} \partial_\mu + g \varepsilon^{abc} \mathcal{A}_\mu^c$ is the total covariant derivative. The ghost energy spectrum is determined by the equation:

$$\det(\nabla^2)^{ab} = 0,$$

where $(\nabla^2)^{ab} = \nabla_\mu^{ac} \nabla^{cb\mu}$. Thus, three branches of the ghost spectrum are obtained $\varepsilon_{gh\,i}^2 = \mathbf{p}^2 + \frac{1}{2} \zeta_i \theta^2$, where $\zeta_1 = -1$, $\zeta_{2,3} = \pm i$, and the ghost contribution to the effective potential is given by the general formula:

$$\mathcal{E}_{gh}^{(1)} = - \sum_{\mathbf{p}, i} \varepsilon_{gh\,i}(\mathbf{p}).$$

Here, the “minus” sign arises as a result of the Fermi statistics applied to ghosts. The branch $\varepsilon_{gh\,1}$ makes a nonzero contribution only to the imaginary part of effective potential. This is easily demonstrated in calculations like those for the gluon contribution to the energy imaginary part. The total result is the following

$$\text{Im } \mathcal{E}_G^{(1)} + \text{Im } \mathcal{E}_{gh}^{(1)} = - \frac{7}{24\sqrt{2}\pi} |\theta|^3.$$

It is also easy to see that the other energy branches make contribution to the real part only

$$\text{Re } \mathcal{E}_{gh}^{(1)reg} = - \mu^{2\epsilon} \sum_{\mathbf{p}, i=2,3} [\mathbf{p}^2 + \frac{1}{2} \zeta_i \theta^2]^{1/2-\epsilon} = - \frac{\mu^{2\epsilon}}{2\pi \Gamma(\epsilon - \frac{1}{2})} \int_0^\infty \frac{ds}{s^{5/2-\epsilon}} \cos \frac{s\theta^2}{2}.$$

Finally, we obtain the following result for the boson contribution to the renormalized effective potential, where only the gluon *physical* degrees of freedom have been retained

$$V_{\text{eff}} \equiv E + \text{Re } \mathcal{E}_G^{(1)} + \text{Re } \mathcal{E}_{\text{gh}}^{(1)} = -\frac{\theta^4}{32g^2} - \frac{|\theta|^3}{2\pi} \left(1 + \frac{3\sqrt{6}}{4}\right).$$

We observe in this connection that the standard prescription for the dimensional regularization integrals, similar to (6), implies that they are assumed to be equal to zero [4]. The terms like $1/\epsilon$ evidently do not appear in the divergent parts. Hence, one may say that the divergent part of the effective potential is put equal to zero by definition, i.e., $\text{Re } \mathcal{E}_G^{(1)\text{div}} = 0$ and $\text{Re } \mathcal{E}_{\text{gh}}^{(1)\text{div}} = 0$, and moreover, masses and coupling constants are not renormalized in this case, which is in agreement with the fact that gauge theories are super-renormalizable in three dimensions [2, 5].

It should also be reminded that the tachyon modes of the gluon spectrum make no contribution to the real part of the correction (4).

We now pass to calculation of the fermion contribution to the effective potential. The quark energy spectrum in the external field specified by (3) can be found from the Dirac equation

$$[(i\partial_\mu\gamma^\mu - m) + \tfrac{1}{2}g\tau^a A_\mu^a\gamma^\mu]\psi(x) = 0. \quad (7)$$

Here, the following representation for the γ -matrices in $2 + 1$ dimensions is chosen: $\gamma^0 = \sigma^3$, $\gamma^{1,2} = i\sigma^{1,2}$, with σ^i as Pauli matrices. The γ -matrices obey the following relation: $\gamma^\mu\gamma^\nu = g^{\mu\nu} - i\varepsilon^{\mu\nu\alpha}\gamma_\alpha$.

Two quantum states of quark in the $SU(2)$ fundamental representation correspond to two different colors. Spinors in $2 + 1$ -dimensions are two-component objects, which corresponds to two degrees of freedom of a fermion (particle–antiparticle) and, hence, spin coefficients are closely related to the sign of the particle energy. The mass term $m\bar{\psi}\psi$ in the Lagrangian is not invariant under $\mathcal{P}$ -inversion of coordinates: $x \rightarrow -x$, $y \rightarrow y$, or $x \rightarrow x$, $y \rightarrow -y$. Recall that simultaneous reflection of two spatial coordinates is equivalent to rotation. The Lagrangian is invariant only under the combined operation that involves $\mathcal{P}$ -inversion and the substitution $m \rightarrow -m$ [1, 6]. Therefore, the spin operator for two-dimensional fermions is naturally introduced in the form $\Sigma = (\text{sgn } m)\gamma^0/2$. Since the matrix γ^0 simultaneously determines the sign of the energy, the positive- and negative-energy states correspond to opposite signs of the spin polarization.

Solving equation (7) in the gauge field (3) results in two different branches of the quark energy spectrum

$$\varepsilon_1^2 = \mathbf{p}^2 + m_{\text{eff } 1}^2, \quad \varepsilon_2^2 = \mathbf{p}^2 + m_{\text{eff } 2}^2,$$

where $m_{\text{eff } 1}^2 = (m - \tilde{\theta})^2$, $m_{\text{eff } 2}^2 = (m - \tilde{\theta})(m + 3\tilde{\theta})$ and $\tilde{\theta} = \theta/4$. It is interesting to note that in the special case, when $m = \tilde{\theta}$, the quark effective mass in this gauge field vanishes, and the two branches of the energy spectrum coincide. For values of the mass lying in the interval $-\tilde{\theta} - 2|\tilde{\theta}| < m < -\tilde{\theta} + 2|\tilde{\theta}|$, the energy squared, ε_2^2 , becomes negative for certain values of the quark momentum $\mathbf{p}$, and tachyonic modes appear in the quark spectrum.

Two branches of the energy spectrum are related to opposite projections of the particle color spin corresponding to two eigenvalues of the operator

$$\mathbf{J} = J^a\tau^a/2 = \tilde{\theta}^{-1}gA^\mu p_\mu - \omega\gamma^0\tau^3(p_\mu\gamma^\mu + \tilde{\theta} - m) \quad (8)$$

that is defined for the plane-wave solutions of the Dirac equation (7) $\psi_s(x) = \exp(-i\varepsilon_s t + i\vec{p}\vec{x})\Psi_s$ with $s = 1, 2$:

$$\mathbf{J}\Psi_s = (-1)^s(\tilde{\theta} - m)\Psi_s,$$

where Ψ_s are constant spinors, corresponding to the Dirac Hamiltonian eigenvalues ε_s . We note, that the r.h.s. of the above equation vanishes in the case $m = \tilde{\theta}$. Moreover, under this condition, it appears that $\Psi_1 = \Psi_2$.

The one-loop quark contribution to the vacuum energy is described by the general equation

$$\mathcal{E}_Q^{(1)} = \text{Re } \mathcal{E}_Q^{(1)} + i \text{Im } \mathcal{E}_Q^{(1)} = - \sum_{\varepsilon_s^2(\mathbf{p}) > 0} \varepsilon_s(\mathbf{p}) - i \sum_{\varepsilon_s^2(\mathbf{p}) < 0} |\varepsilon_s(\mathbf{p})|. \quad (9)$$

The real part of the correction is defined by the expression

$$\text{Re } \mathcal{E}_Q^{(1)} = -\frac{1}{(2\pi)^2} \int d\mathbf{p} \varepsilon_1 - \frac{1}{(2\pi)^2} \int_{|\mathbf{p}|=p_*}^{\infty} d\mathbf{p} \varepsilon_2.$$

Here $p_* = [-m_{\text{eff}2}^2]^{1/2} \Theta(-m_{\text{eff}2}^2)$ is the lowest of the quark momentum values that provide a real contribution to the quark energy in the absence of the tachyonic modes. Function $\Theta(x)$ is the Heaviside (unit step) function. To find the finite part of the quark energy correction, the same procedure as in the case of gluons can be applied. The negative sign in front of the real part of the energy in (9) is due to the fact that quarks are fermions. The quark contribution to the effective potential finally reads

$$\begin{aligned} \text{Re } \mathcal{E}_Q^{(1)} = \frac{1}{6\pi} \Big\{ & |m - \tilde{\theta}| \left((m - \tilde{\theta})^2 - \frac{3|m|}{4} |m + 3\tilde{\theta}| \right) - \frac{|m|}{4} (m - \tilde{\theta})(5m - \tilde{\theta}) + \\ & |m| \tilde{\theta}^2 + [(m - \tilde{\theta})(m + 3\tilde{\theta})]^{3/2} \Theta((m + \tilde{\theta})^2 - 4\tilde{\theta}^2) \Big\}. \end{aligned}$$

We note that the sign of this expression is opposite to that of the gluon contribution.

Like in the gluon case, the imaginary part of the energy does not diverge and takes nonzero values only if tachyonic modes are present. The final expression for the imaginary part can be written in the following simple form

$$\text{Im } \mathcal{E}_Q^{(1)} = -\frac{1}{6\pi} |(m - \tilde{\theta})(m + 3\tilde{\theta})|^{3/2} \Theta(4\tilde{\theta}^2 - (m + \tilde{\theta})^2).$$

In particular, in the limit of the vanishing quark mass, $m \rightarrow 0$, we obtain

$$\text{Re } \mathcal{E}_Q^{(1)} = \frac{|\tilde{\theta}|^3}{6\pi}, \quad \text{Im } \mathcal{E}_Q^{(1)} = -\frac{\sqrt{3}|\tilde{\theta}|^3}{2\pi}.$$

It should be emphasized that it is the tachyonic modes in the quark spectrum that lead to the appearance of an imaginary part in the energy of vacuum fluctuations. This means that the constant gauge field configuration in question is unstable. However, such factors, as boundaries and finite volume of the spatial region occupied by the gauge field, finite temperature, as well as consideration for inhomogeneous modes and higher loop contributions may lead to its stabilization.

References

- [1] *Deser S., Jackiw R., Templeton S.* Ann. Phys. (N.Y.) **1982**. Vol. 140. p. 372.
- [2] *Jackiw R., Templeton S.* Phys. Rev. **1981**. Vol. D23. p. 2291.
- [3] *Sokolov A.A. et al.* Kalibrovochnye polya (Gauge Fields), Moscow: Moscow University Press, 1986, [in Russian].
- [4] *Faddeev L.D., Slavnov A.A.* Gauge Fields: Introduction to Quantum Theory, Benjamin-Cummings Publishing Co., 1980. *Trottier H.D.* Phys. Rev. **1991**. Vol. D44. p. 464.
- [5] *Templeton S.* Phys. Rev. **1981**. Vol. D24. p. 3134.
- [6] *Deser S., Jackiw R., Templeton S.* Phys. Rev. Lett. **1982**. Vol. 48. p. 975.